



IUB™

Ballerine® Update:


Global Vigilance Data Q2/20*

IUB™ Ballerine® MIDI

Total Estimated Insertions since Q1 2015: **94,962**

Event	Rate in %	
Expulsion	3.57%	
Pregnancy	0.48%	
Ectopic Pregnancy	0.01%	>90% LESS RISK*
Perforation	0.07%	39% LESS RISK*

*compared to T-IUDs




T-IUD:
5%³



T-Cu IUD:
1%³



T-IUD:
>0.26%^{4,5}



T-IUD:
1:909²

1 based on accumulated data from Q1 2015 – Q2 2020 on total estimated insertions and OCON's ongoing vigilance reporting

2 Heinemann et al. Risk of Uterine Perforation with Levonorgestrel-Releasing and Copper Intrauterine Devices in the European. Active Surveillance Study on Intrauterine Devices. Contraception (2015), doi: 10.1016/j.contraception.2015.01.007.

3 Aoun et al. Effects of age, parity, and device type on complications and discontinuation of intrauterine devices. Obstetrics & Gynecology (123:3) 2014, 585-592.

4 <https://www.sciencedirect.com/science/article/pii/S0010782415000128>

5 [https://www.fertstert.org/article/S0015-0282\(00\)01605-8/pdf](https://www.fertstert.org/article/S0015-0282(00)01605-8/pdf)